

**Bioavailability of pollutants and chemotaxis**

Tino Krell<sup>1</sup>, Jesús Lacal<sup>1</sup>, Jose Antonio Reyes-Darias<sup>1</sup>, Celia Jimenez-Sanchez<sup>2</sup>,  
Rungroch Sungthong<sup>2</sup> and Jose Julio Ortega-Calvo<sup>2\*</sup>

<sup>1</sup> Department of Environmental Protection, CSIC, Estación Experimental del Zaidín,  
Profesor Albareda 1, 18008-Granada, Spain

<sup>2</sup> Instituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS-CSIC), Apartado  
1052, 41080-Seville, Spain

\*Corresponding author: tel: (+34) 95-4624711; fax: (+34) 95-4624002;  
e-mail: jjortega@irnase.csic.es

## **Abstract**

The exposure of bacteria to pollutants induces frequently chemoattraction or chemorepellent reactions. Recent research suggests that the capacity to degrade a toxic compound has co-evolved in some bacteria with the capacity to chemotactically react to it. There is an increasing amount of data which show that chemoattraction to biodegradable pollutants increases their bioavailability which translates into an enhancement of the biodegradation rate. Pollutant chemoreceptors so far identified are encoded on degradation or resistance plasmids. Genetic engineering of bacteria, such as the transfer of chemoreceptor genes, offers thus the possibility to optimize biodegradation processes.

## **Introduction**

Biodegradation of anthropogenic organic chemicals (AOCs) in natural and engineered environments is often not as efficient as expected due to a limited bioavailability, which represents the accessibility of a chemical for biotransformation and toxicity. As a result of sorption to soils and sediments, pollutants often only exhibit weak chemical activity gradients that promote their uptake and transformation by cells. Thus, the biodegradation rates may reflect the dependencies of restricted phase exchanges, and the pollutants, together with their environmental risks, may persist for longer periods of time (Figure 1). Research over the last decade has shown that the chemotactic movement of bacteria increases bioavailability, which in turn was found to have a beneficial role in bioremediation. In this review, we consider chemotaxis as the diverse tactic reactions to pollutants of bacteria exhibiting flagellar motility. The focus will be on AOCs, but due to their environmental relevance, the recent advances in the field of chemotaxis to inorganic pollutants, such as metals and nanomaterials, will also be reviewed.

## **Chemotaxis towards and away from pollutants**

Chemotaxis has been extensively studied in enterobacteria that show chemotaxis to a limited number of compounds like amino acids, organic acids and sugars [1]. Many free-living bacteria have an increased number of chemoreceptors, which were shown to mediate chemotaxis to a wider range of compounds as compared to enterobacteria [2,3]. Interestingly, many of these compounds are chemicals of environmental concern. Chemoattraction was observed for example towards biphenyl, benzoic acid and chlorobenzoic acids [4\*\*], toluene and its derivatives [5,6\*], naphthalene and its derivatives [6\*,7], nitroaromatics [8], chloroaromatics [9], chloronitroaromatics [10\*],

1 aminoaromatics [11], explosives [12], aliphatic hydrocarbons [13] and herbicides [14\*]  
2 in species like *Rhizobium* sp., *Bradyrhizobium* sp., *Pseudomonas* sp., *Azospirillum* sp.,  
3 *Ralstonia* sp., *Burkholderia* sp. or *Flavimonasoryzihabitans*. In a significant number of  
4 cases the physiological relevance of chemoattraction to pollutants lies in the fact that  
5 these compounds serve as carbon and energy sources. This may be exemplified by the  
6 chemotaxis towards toluene and naphthalene by *Pseudomonas putida* DOT-T1E and  
7 G7, which possess specific degradation routes for both compounds, respectively  
8 [15,16]. In some cases chemoattraction was observed towards pollutants that are not  
9 metabolized by the bacterium and the physiological relevance of this behavior is little  
10 understood.

11  
12 Given the toxic potential of most pollutants, it is conceivable that bacteria have also  
13 evolved chemorepellent responses. Bacterial repellence has been reported, for example,  
14 to hydrogen peroxide, hypochlorite and N-chlorotaurine [17], the PAHs anthracene and  
15 pyrene [18],  $\text{Co}^{++}$  and  $\text{Ni}^{++}$  [19], and silver nanoparticles [20]. Some chemicals can even  
16 be chemo-attractants for one bacterial species and be repellent for another [5,21,22].  
17 The physical state of the chemical also appears to influence the type of response, since it  
18 was shown that the naphthalene degrader *P. putida* G7 was repelled by naphthalene in  
19 the vapor phase, whereas it was attracted when the compound was dissolved in the  
20 aqueous phase [23\*\*]. In the light of such results one has to keep in mind that an  
21 observed chemotaxis phenotype can be the result of the action of several, potential  
22 antagonistic chemoreceptors that differ in their sensitivity to a given compound.

23  
24 The complexity of the chemotactic reactions to pollutants is also illustrated by the  
25 dissimilar reactions exhibited by *P. putida* to silver nanoparticles and  $\text{Ag}^+$  ions [20].

Nano-scale silver induced a repellent response, possibly due to a direct effect of the nanoparticles on bacterial cells, and not due to the release of soluble  $\text{Ag}^+$  from the particles. Indeed, the bacterium did not show any repellent response to soluble  $\text{Ag}^+$ , what is in agreement with earlier reports, which demonstrate that some harmful chemicals, such as  $\text{Cu}^{++}$ , do not induce negative taxis in bacteria [19]. Furthermore a positive tactic response was detected at low concentrations of silver nitrate, what probably reflects the physiological role of  $\text{Ag}^+$  ions and other metal ions like  $\text{Mn}^{3+}$  and  $\text{Fe}^{3+}$  towards which taxis was observed [24-26]. The bacterial repellent responses can be considered as a prelude of toxicity because it is often observed only at sub-lethal pollutant concentrations, which suggests that taxis may indeed be part of survival strategies aimed on minimizing the deleterious effects of toxic compounds. In addition, the molecular machinery for the detection of chemicals for tactic purposes can also be employed for analytical purposes, namely the development of alternative bioassay methods.

#### **Pollutant chemoreceptors so far identified are present on plasmids**

The specificity of a chemotactic response is determined by chemoreceptors. Two chemoreceptors for aromatic pollutants have so far been described, which are NahY [27] of the naphthalene degrading *P. putida* G7 and McpT of the toluene, benzene and ethylbenzene degrading *P. putida* DOT-T1E [6\*]. Both receptors mediate chemoattraction towards their respective degradation substrates. Interesting parallels exist between both receptors. NahY and McpT are encoded on plasmids pNAH7 [28] and pGRT1 [29], respectively. Both plasmids contain genes that are related to either degradation of or resistance to aromatic pollutants. The pNAH7 plasmid contains genes that encode the naphthalene degradation route and the *nahY* gene is co-transcribed with

part of these genes [27]. Plasmid pGRT1 contains two *mcpT* alleles which are both in vicinity of the *ttgGHI* operon that encodes the primary efflux pump responsible for solvent resistance [30]. McpT was found to mediate an extreme form of chemoattraction, termed hyperchemotaxis, towards a wide range of mono- and biaromatic compounds [6\*]. The capacity of McpT to mediate a hyperchemotaxis response towards crude oil samples is illustrated in Figure 2. Due to taxis and a very high solvent resistance, cells were able to assemble on the surface of this toxic mixture of compounds. The deletion of the *mcpT* gene (Figure 2) abolished this capacity.

Although NahY and McpT exert a similar function, the sequence alignment of their ligand binding regions (LBRs) reveals no significant identity. However, close homologues of both receptors are found on other degradation plasmids. For example receptors showing 99 % sequence identity with McpT are found on the carbazole degradation plasmids pCAR1 of *P. resinovorans* CA10 [31] or the toluene degradation plasmid pWW53 of *P. putida* MT53 [32]. In analogy, NahY homologues are found on the naphthalene degradation plasmids pDTG1 [33] or pND6-1 [34]. This suggests that there are at least two different families of pollutant chemoreceptors, of which NahY and McpT are representative members.

A chemosensory signaling cascade is formed by chemoreceptors and cytosolic signaling proteins [1\*]. Plasmids mentioned contain chemoreceptor genes but lack those of signaling proteins. This implies that chemotaxis is mediated by the concerted action of plasmid encoded receptors and genome encoded signaling proteins. A transfer of the *mcpT* gene into strains *P. putida* KT2440 and F1 conferred the hyperchemotaxis phenotype to both strains [6\*]. This indicates that McpT is able to interact with the

1 signaling proteins, which offers the possibility of conferring pollutant chemotaxis to  
2 other bacteria by chemoreceptor gene transfer. Although these pollutant receptors are  
3 plasmid-encoded there is also evidence for genome-encoded pollutant chemoreceptors,  
4 since for example the plasmid-free strains *P. putida* KT2440 or F1 show chemotaxis  
5 towards toluene which, however, is in its magnitude inferior to the McpT-mediated  
6 hyperchemotaxis [6\*].

## 7

### 8 **Chemoattraction increases bioavailability of pollutants and enhances** 9 **biodegradation rate**

10 There is now sufficient evidence demonstrating that chemoattraction increases the  
11 bioavailability of pollutants. The best studied example is the capacity of *P. putida* G7 to  
12 degrade naphthalene. Grimm and Harwood [27] have proposed that NahY-mediated  
13 taxis towards naphthalene might facilitate its biodegradation. Proof of this hypothesis  
14 was brought by Aitken and co-workers. Using a heterogeneous aqueous system they  
15 were able to demonstrate that chemotaxis enhances naphthalene biodegradation [35].  
16 Subsequent studies using chemotactic and non-chemotactic strains of *P. putida* G7  
17 clearly demonstrated that chemotaxis increased naphthalene degradation when the  
18 compound is present in a non-aqueous-phase liquid [36]. Bioavailability can also be  
19 promoted by the chemotactic transport of *P. putida* G7 through fungal mycelia that act  
20 as pathways for mobilization [37\*]. There are several studies that compare the capacity  
21 of microorganisms to degrade different pollutants with their capacity to chemotactically  
22 approach these compounds. Interestingly, in some cases chemotaxis was only observed  
23 towards compounds which were degraded by the microorganism whereas structurally  
24 similar non-substrate compounds were not found to be chemoattractants [10\*,38],  
25 which confirms the link between chemotaxis and biodegradation.

1 The standard chemotaxis assays involve the measurement of motility in buffer  
2 solutions. However, pollutants adsorb frequently onto solids or into pores. To evaluate  
3 whether efficient pollutant chemotaxis occurs also under these more *in situ* conditions,  
4 chemotaxis assays were developed to monitor taxis in contaminated soils. Research  
5 based on close to *in situ* conditions has shown that pollutant chemotaxis occurs also in  
6 contaminated soil [39\*\*] and in a reconstituted bench-scale microcosm [40\*\*].  
7 Biodegradation studies of carbon tetrachloride [41] and pesticides [42] have also  
8 demonstrated the ability of chemotaxis to enhance biodegradation under laboratory-  
9 conditions. It has also been shown in later research that bacterial motility and transport  
10 can be controlled through a suitable choice of chemical effectors [20,43,44\*]. In well-  
11 controlled column systems, the influence of different effectors on the deposition of *P.*  
12 *putida* G7 was assessed in selected porous environments. Cellular deposition, however,  
13 was concomitantly dependent on the cellular motility (hyper-motility, attraction or  
14 repulsion), the sorption of the effector to the column packing material, and the resulting  
15 pore-water concentration (Figure 3).

16  
17 **Conclusions: Genetic engineering to improve resistance, degradation pathway**  
18 **expression and chemotactic mobilization.**

19 The use of genetically engineered microorganisms for biodegradation purposes has been  
20 crowned with little success in the past. However, the reasons for the reduced efficiency  
21 were clearly identified and include, amongst others, limited bacterial resistance towards  
22 toxic compounds, inadequate expression of degradation pathways and bioavailability  
23 restrictions [45]. Research over the last decade has advanced enormously the  
24 understanding of the basic mechanisms related to these issues. The new knowledge on  
25 the functional link between chemotaxis and biodegradation will have implications for



the design of new bioaugmentation and biostimulation strategies oriented toward improving the performance of pollutant-degrading microbial populations. These strategies may overcome, for example, the limitations usually caused by the use of inoculants designed exclusively on the basis of their catabolic potential. In parallel, major conceptual and technical advances have been made in the field of synthetic biology, which now permit a much more rational and directed modification of the bacterial genome [46]. In the case of chemotaxis to pollutants it may be possible to engineer the process by transferring the chemoreceptor gene which establishes a signaling cascade with host signaling genes. Genetic engineering combining the acquired knowledge of pollutant related chemical processes with powerful synthetic biology approaches should give rise to biodegrading microorganisms that perform better than their predecessors.

## **Acknowledgements**

During the preparation of the manuscript, our research laboratories at EEZ-CSIC and IRNAS-CSIC were supported by the Spanish Ministry of Economy and Competitiveness (grants Bio2010-16937 and CGL2010-22068-C02-01), Junta de Andalucía (P09-RNM-4509 and RNM-312) and the BBVA Foundation (BIOCON08 185/09).

## **References and recommended reading**

Papers of particular interest have been highlighted as:

\* of special interest

\*\* of outstanding interest

1. Baker MD, Wolanin PM, Stock JB: **Signal transduction in bacterial chemotaxis.** *Bioessays* 2006, **28**:9-22.
2. Krell T, Lacal J, Munoz-Martinez F, Reyes-Darias JA, Cadirci BH, Garcia-Fontana C, Ramos JL: **Diversity at its best: bacterial taxis.** *Environ Microbiol* 2011, **13**:1115-1124.
3. Lacal J, Garcia-Fontana C, Munoz-Martinez F, Ramos JL, Krell T: **Sensing of environmental signals: classification of chemoreceptors according to the size of their ligand binding regions.** *Environ Microbiol* 2010, **12**:2873-2884.
4. Tremaroli V, Vacchi Suzzi C, Fedi S, Ceri H, Zannoni D, Turner RJ: **Tolerance of *Pseudomonas pseudoalcaligenes* KF707 to metals, polychlorobiphenyls and chlorobenzoates: effects on chemotaxis-, biofilm- and planktonic-grown cells.** *FEMS Microbiol Ecol* 2010, **74**:291-301.
- \*\* First investigation of co-influence of metals and polychlorobiphenyls on bacterial chemotaxis resulted that different types of metals could affect the chemotactic response of bacteria. The article also emphasized the connection between bacterial chemotaxis and biofilm formation in such condition with the presence of both pollutant and specific metal that may be an effective parameter to improve bioremediation.
5. Parales RE, Ditty JL, Harwood CS: **Toluene-degrading bacteria are chemotactic towards the environmental pollutants benzene, toluene, and trichloroethylene.** *Appl Environ Microbiol* 2000, **66**:4098-4104.
6. Lacal J, Munoz-Martinez F, Reyes-Darias JA, Duque E, Matilla M, Segura A, Ortega-Calvo JJ, Jimenez-Sanchez C, Krell T, Ramos JL: **Bacterial chemotaxis towards aromatic hydrocarbons in *Pseudomonas*.** *Environ Microbiol* 2011, **13**:1733-1744.

- \* Well-defined regulatory gene expression of the chemotactic phenotype was depended on dosing of chemical effectors tested.
7. Grimm AC, Harwood CS: **Chemotaxis of *Pseudomonas* spp. to the polyaromatic hydrocarbon naphthalene.** *Appl Environ Microbiol* 1997, **63**:4111-4115.
8. Iwaki H, Muraki T, Ishihara S, Hasegawa Y, Rankin KN, Sulea T, Boyd J, Lau PC: **Characterization of a pseudomonad 2-nitrobenzoate nitroreductase and its catabolic pathway-associated 2-hydroxylaminobenzoate mutase and a chemoreceptor involved in 2-nitrobenzoate chemotaxis.** *J Bacteriol* 2007, **189**:3502-3514.
9. Gordillo F, Chavez FP, Jerez CA: **Motility and chemotaxis of *Pseudomonas* sp. B4 towards polychlorobiphenyls and chlorobenzoates.** *FEMS Microbiol Ecol* 2007, **60**:322-328.
10. Pandey J, Sharma NK, Khan F, Ghosh A, Oakeshott JG, Jain RK, Pandey G: **Chemotaxis of *Burkholderia* sp. Strain SJ98 towards chloronitroaromatic compounds that it can metabolise.** *BMC Microbiol* 2012, **12**:19. doi: 10.1186/1471-2180-12-19
- \* The link of chemotaxis with biodegradation potential was emphasized. Interestingly, analogue compounds that were not used for growth did not induce chemotaxis in the bacterium tested.
11. Parales RE: **Nitrobenzoates and aminobenzoates are chemoattractants for *Pseudomonas* strains.** *Appl Environ Microbiol* 2004, **70**:285-292.
12. Leungsakul T, Keenan BG, Smets BF, Wood TK: **TNT and nitroaromatic compounds are chemoattractants for *Burkholderia cepacia* R34 and *Burkholderia* sp. strain DNT.** *Appl Microbiol Biotechnol* 2005, **69**:321-325.

13. Lanfranconi MP, Alvarez HM, Studdert CA: **A strain isolated from gas oil-contaminated soil displays chemotaxis towards gas oil and hexadecane.** *Environ Microbiol* 2003, **5**:1002-1008.
14. Liu X, Parales RE: **Bacterial chemotaxis to atrazine and related s-triazines.** *Appl Environ Microbiol* 2009, **75**:5481-5488.
- \* A well-defined example of relationships between chemotactic response and pollutant catabolism in bacteria with a case study of the triazine herbicide atrazine. Interestingly, both metabolized and non-metabolized pollutants were attractants for the bacteria tested.
15. Mosqueda G, Ramos-Gonzalez MI, Ramos JL: **Toluene metabolism by the solvent-tolerant *Pseudomonas putida* DOT-T1 strain, and its role in solvent impermeabilization.** *Gene* 1999, **232**:69-76.
16. Tsuda M, Iino T: **Naphthalene degrading genes on plasmid NAH7 are on a defective transposon.** *Mol Gen Genet* 1990, **223**:33-39.
17. Benov L, Fridovich I: ***Escherichia coli* exhibits negative chemotaxis in gradients of hydrogen peroxide, hypochlorite, and N-chlorotaurine: Products of the respiratory bursts of phagocytic cells.** *Proc Natl Acad Sci USA* 1996, **93**:4999-5002.
18. Ortega-Calvo JJ, Marchenko AI, Vorobyov AV, Borovick RV: **Chemotaxis in polycyclic aromatic hydrocarbon-degrading bacteria isolated from coal-tar- and oil-polluted rhizospheres.** *FEMS Microbiology Ecology* 2003, **44**:373-381.
19. Tso W-W, Adler J: **Negative chemotaxis in *Escherichia coli*.** *J Bacteriol* 1974, **118**:560-576.

20. Ortega-Calvo JJ, Molina R, Jimenez-Sanchez C, Dobson PJ, Thompson IP:  
**Bacterial tactic response to silver nanoparticles.** *Environ Microbiol Rep* 2011,  
**3**:526-534.
21. Vardar G, Barbieri P, Wood TK: **Chemotaxis of *Pseudomonas stutzeri* OX1 and  
*Burkholderia cepacia* G4 toward chlorinated ethenes.** *Appl Microbiol Biotechnol*  
2005, **66**:696-701.
22. Shitashiro M, Tanaka H, Hong CS, Kuroda A, Takiguchi N, Ohtake H, Kato J:  
**Identification of chemosensory proteins for trichloroethylene in *Pseudomonas*  
*aeruginosa*.** *J Biosci Bioeng* 2005, **99**:396-402.
23. Hanzel J, Harms H, Wick LY: **Bacterial chemotaxis along vapor-phase gradients  
of naphthalene.** *Environ Sci Technol* 2010, **44**:9304-9310.
- \*\* The article demonstrated the influence of physicochemical properties of the effector  
on chemotactic response in bacteria. Naphthalene causes in the aqueous phase  
chemoattraction by *Pseudomonas putida* PpG7, but, when the chemical is present in  
vapor phase, it acts as a repellent, even at lower doses than in the aqueous phase.
24. de Sanchez SR, Schiffrin DJ: **Bacterial chemo-attractant properties of metal ions  
from dissolving electrode surfaces.** *J Electroanal Chem* 1996, **403**:39-45.
25. Bencharit S, Ward MJ: **Chemotactic responses to metals and anaerobic electron  
acceptors in *Shewanella oneidensis* MR-1.** *J Bacteriol* 2005, **187**:5049-5053.
26. Childers SE, Ciufo S, Lovley DR: ***Geobacter metallireducens* accesses insoluble  
Fe(III) oxide by chemotaxis.** *Nature* 2002, **416**:767-769.
27. Grimm AC, Harwood CS: **NahY, a catabolic plasmid-encoded receptor required  
for chemotaxis of *Pseudomonas putida* to the aromatic hydrocarbon  
naphthalene.** *J Bacteriol* 1999, **181**:3310-3316.

28. Sota M, Yano H, Ono A, Miyazaki R, Ishii H, Genka H, Top EM, Tsuda M:  
**Genomic and functional analysis of the IncP-9 naphthalene-catabolic plasmid  
NAH7 and its transposon Tn4655 suggests catabolic gene spread by a tyrosine  
recombinase.** *J Bacteriol* 2006, **188**:4057-4067.
29. Molina L, Duque E, Gomez MJ, Krell T, Lacal J, Garcia-Puente A, Garcia V,  
Matilla MA, Ramos JL, Segura A: **The pGRT1 plasmid of *Pseudomonas putida*  
DOT-T1E encodes functions relevant for survival under harsh conditions in the  
environment.** *Environ Microbiol* 2011, **13**:2315-2327.
30. Rojas A, Duque E, Mosqueda G, Golden G, Hurtado A, Ramos JL, Segura A:  
**Three efflux pumps are required to provide efficient tolerance to toluene in  
*Pseudomonas putida* DOT-T1E.** *J Bacteriol* 2001, **183**:3967-3973.
31. Miyakoshi M, Shintani M, Terabayashi T, Kai S, Yamane H, Nojiri H:  
**Transcriptome analysis of *Pseudomonas putida* KT2440 harboring the  
completely sequenced IncP-7 plasmid pCAR1.** *J Bacteriol* 2007, **189**:6849-6860.
32. Yano H, Garruto CE, Sota M, Ohtsubo Y, Nagata Y, Zylstra GJ, Williams PA,  
Tsuda M: **Complete sequence determination combined with analysis of  
transposition/site-specific recombination events to explain genetic organization  
of IncP-7 TOL plasmid pWW53 and related mobile genetic elements.** *J Mol Biol*  
2007, **369**:11-26.
33. Dennis JJ, Zylstra GJ: **Complete sequence and genetic organization of pDTG1,  
the 83 kilobase naphthalene degradation plasmid from *Pseudomonas putida*  
strain NCIB 9816-4.** *J Mol Biol* 2004, **341**:753-768.
34. Li W, Shi J, Wang X, Han Y, Tong W, Ma L, Liu B, Cai B: **Complete nucleotide  
sequence and organization of the naphthalene catabolic plasmid pND6-1 from  
*Pseudomonas* sp. strain ND6.** *Gene* 2004, **336**:231-240.

35. Marx RB, Aitken MD: **Bacterial chemotaxis enhances naphthalene degradation in a heterogeneous system.** *Environ Sci Technol* 2000, **34**:3379-3383.
36. Law AM, Aitken MD: **Bacterial chemotaxis to naphthalene desorbing from a nonaqueous liquid.** *Appl Environ Microbiol* 2003, **69**:5968-5973.
37. Furuno S, Pazolt K, Rabe C, Neu TR, Harms H, Wick LY: **Fungal mycelia allow chemotactic dispersal of polycyclic aromatic hydrocarbon-degrading bacteria in water-unsaturated systems.** *Environ Microbiol* 2010, **12**:1391-1398.
- \* The study evidenced that bacterial chemotaxis requires a liquid phase for motility. Interestingly, the liquid films surrounding fungal mycelia provide appropriate environments for chemical gradients and chemotactic responses by motile bacteria.
38. Samanta SK, Bhushan B, Chauhan A, Jain RK: **Chemotaxis of a *Ralstonia* sp. SJ98 toward different nitroaromatic compounds and their degradation.** *Biochem Biophys Res Commun* 2000, **269**:117-123.
39. Paul D, Singh R, Jain RK: **Chemotaxis of *Ralstonia* sp. SJ98 towards p-nitrophenol in soil.** *Environ Microbiol* 2006, **8**:1797-1804.
- \*\* Tracking of bacterial dispersal in heterogeneous soil with various qualitative and quantitative methods. The study shows chemotactic bacteria as valuable agents for soil bioremediation.
40. Strobel KL, McGowan S, Bauer RD, Griebler C, Liu J, Ford RM: **Chemotaxis increases vertical migration and apparent transverse dispersion of bacteria in a bench-scale microcosm.** *Biotechnol Bioeng* 2011, **108**:2070-2077.
- \*\* Experimental model of great value for studying bacterial chemotaxis and bacteria-pollutant interaction in sand aquifers and *in situ* bioremediation.

41. Witt ME, Dybas MJ, Worden RM, Criddle CS: **Motility-enhanced bioremediation of carbon tetrachloride-contaminated aquifer sediments.** *Environ Sci Technol* 1999, **33**:2958-2964.
42. Hawkins AC, Harwood CS: **Chemotaxis of *Ralstonia eutropha* JMP134(pJP4) to the herbicide 2,4-dichlorophenoxyacetate.** *Appl Environ Microbiol* 2002, **68**:968-972.
43. Velasco-Casal P, Wick LY, Ortega-Calvo JJ: **Chemoeffectors decrease the deposition of chemotactic bacteria during transport in porous media.** *Environ Sci Technol* 2008, **42**:1131-1137.
44. Jimenez-Sanchez J, Wick LY, Ortega-Calvo JJ: **Chemical effectors cause different motile behavior and deposition of bacteria in porous media.** *Environ Sci Technol* 2012, **46**:6790-6797.
- \* The paper provides new information about the interactions on chemotactic bacteria and porous materials. The results contained provide clear evidences for a modulation of bacterial transport in saturated porous media depending on chemical effectors.
45. Cases I, de Lorenzo V: **Genetically modified organisms for the environment: stories of success and failure and what we have learned from them.** *Int Microbiol* 2005, **8**:213-222.
46. Cambray G, Mutalik VK, Arkin AP: **Toward rational design of bacterial genomes.** *Curr Opin Microbiol* 2011, **14**:624-630.



## Figure Legends

**Figure 1) Conceptual framework of this review.** Biodegradation, bioavailability and chemotaxis are interconnected through a variety of concepts related to bacterial physiology and genetics, bioremediation performance and environmental risk.

**Figure 2) Chemotaxis of *P. putida* DOT-T1E (A) and its mutant devoid of *mcpT* genes (B) towards undiluted crude oil recovered from the Spanish coast following the “Prestige” oil tanker accident.** In the wild type strain cells accumulate right on the surface of this highly toxic mixture of hydrocarbons. Mutation of the *mcpT* gene abolished taxis. Reproduced with permission from [6\*].

**Figure 3) Effect of exposure of *Pseudomonas putida* G7 to salicylate (promoting positive taxis) and to silver nanoparticles (repellence) on bacterial transport through sand (A) and motile behavior (B-D).** Salicylate significantly reduced deposition of G7 cells, whereas AgNPs enhanced attachment and caused filter blocking that resulted in a progressive decrease in deposition (A). Computer-aided analysis of cell trajectories showed (C-D) that exposure to salicylate induced smooth cell movement with no turning events (peaks in the rate of change of direction or RCDI), whereas cells exposed to AgNPs exhibited tortuous movement. Modified with permission from [44\*].





FIGURE 1. Krell et al.

*Pseudomonas putida* DOT-T1E



*Pseudomonas putida* DOT-T1E  $\Delta mcpT$




FIGURE 2. Krell et al.




FIGURE 3. Krell et al.